

2014-2015 School Year

Below are the number of doses and each vaccine required for school entry. Changes from the 2013-2014 requirements are indicated in **blue**.

3 to 5 years old	3 Hep B (Hepatitis B)	
	4 DTaP (Diphtheria, Tetanus & Pertussis)	
	3 Polio (Inactivated Polio)	
	1 MMR (Measles, Mumps & Rubella)	
	1 Varicella	
Kindergarten	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	2 Hep A (Hepatitis A)
Grades 1 to 5	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	
Grades 6 to 11	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	1 Tdap (Tetanus & Pertussis)
		1 MCV4 (Meningococcal)
Grade 12	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	1 Tdap (Tetanus & Pertussis)
		2 MCV4 (Meningococcal)

Hep B The minimum age for the 3rd dose of Hepatitis B is 24 weeks of age.

DTaP Four doses of DTaP/DTP/DT are acceptable if 4th dose was administered on or after child's 4th birthday.

Polio Three doses of Polio are acceptable for all grade levels if the third dose was given on or after the 4th birthday and at least 6 months after the previous dose with only one type of vaccine used (all OPV or all IPV).

For students in grades kindergarten through 4th grade, the final dose must be administered on or after the 4th birthday, and be administered **at least 6 months** after the previous dose.

Live Vaccines (MMR, Varicella & LAIV) Live vaccines that are not administered on the same day must be administered a minimum of 28 days apart. The second dose should be repeated if the doses are separated by less than 28 days.

Varicella Physician documentation of disease history, including month and year, is proof of immunity for children entering preschool through 6th grade. Parental report of disease history is acceptable for grades 7-12.

Tdap There is no minimum interval from the last Td dose.

MCV4 Individuals who receive dose 1 after their 16th birthday only need 1 dose of MCV4.

2015-2016 School Year (Proposed)

Below are the number of doses and each vaccine required for school entry. Changes from the 2014-2015 requirements are indicated in **blue**.

3 to 5 years old	3 Hep B (Hepatitis B)	
	4 DTaP (Diphtheria, Tetanus & Pertussis)	
	3 Polio (Inactivated Polio)	
	1 MMR (Measles, Mumps & Rubella)	
	1 Varicella	
K -1st grade	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	2 Hep A (Hepatitis A)
Grades 2 to 5	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	
Grades 6 to 11	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	1 Tdap (Tetanus & Pertussis)
		1 MCV4 (Meningococcal)
Grade 12	3 Hep B	2 MMR
	5 DTaP	2 Varicella
	4 Polio	1 Tdap (Tetanus & Pertussis)
		2 MCV4 (Meningococcal)

Hep B The minimum age for the 3rd dose of Hepatitis B is 24 weeks of age.

DTaP Four doses of DTaP/DTP/DT are acceptable if 4th dose was administered on or after child's 4th birthday.

Polio Three doses of Polio are acceptable for all grade levels if the third dose was given on or after the 4th birthday and at least 6 months after the previous dose with only one type of vaccine used (all OPV or all IPV).

For students in grades kindergarten through 4th grade, the final dose must be administered on or after the 4th birthday, and be administered **at least 6 months** after the previous dose.

Live Vaccines (MMR, Varicella & LAIV) Live vaccines that are not administered on the same day must be administered a minimum of 28 days apart. The second dose should be repeated if the doses are separated by less than 28 days.

Varicella Physician documentation of disease history, including month and year, is proof of immunity for children entering preschool through 7th grade. Parental report of disease history is acceptable for grades 8-12.

Tdap There is no minimum interval from the last Td dose.

MCV4 Individuals who receive dose 1 on or after their 16th birthday only need 1 dose of MCV4.